



PROBABILITY

UNIT-III

Introduction:— Galileo (1564-1624) of an Italian Mathematician was first to attempt a quantitative measure of Probability while dealing with some Problems related to the theory of dice in Gambling. For the first foundation of mathematical theory Probability was laid in middle of 17th Century by the French Mathematician Pascal and Fermat while solving no. of Problems of French Gamblers.

Russian mathematicians are also have made a very valuable contribution on modern theory of Probability. Famous mathematical scientist Chebyshev's who founded Russian School of Mathematics and Statistics. They are proposed on definitions on Probability.

Random experiment:— Consider an experiment can throw repeated ascetically identical conditions, does not gives unique results. But result is any one of the several possible outcomes. This type of an experiment is called random experiment.

Ex:— Tossing a coin, rolling a dice... etc.



Sample Space :- Set of all possible outcomes of an experiment is known as sample space. It is denoted with S or Ω

Ex:- ① n coins are ~~thrown~~ ^{rolled} sample space is 2^n

② n dice are rolled sample space is 6^n .


③ In a pack of cards if we draw x cards then sample space is ${}^{52}C_x$.

④ In an urn contains m -white n -blue k -yellow balls if we draw x balls then sample space ${}^{m+n+k}C_x$.

Event :- the outcome of an experiment is known as an event.

Ex:- ① Throwing a coin and getting a head is an event.

② Rolling a dice and getting 5th face is an event.



Simple event :- An event is said to be simple event if it has single possible ~~out~~ ^{outcome}.

Ex:- Rolling a dice getting sixth face.



Compound event:— An event is said to be compound event if it can be split into two.

Ex: Rolling a dice getting sum is 9.
 $S = \{(6,3) (3,6) (5,4) (4,5)\}$.

Exhaustive Cases:— The total number of possible outcomes in an experiment is known as exhaustive cases.

Ex: (1) In tossing a coin total exhaustive cases are 2^1
(2) In rolling a dice total exhaustive cases are 6^1

Favourable Cases:— The number of cases favourable to an event in a trial is known as favourable cases.

Mutually exclusive cases events:— The events are said to be mutually exclusive if happening of one event prevents the happening of another event is called mutually exclusive events.

Independent events:— The events are said to be independent events if happening of one event has no influence of happening of other event are said to be independent events.



Mathematical (or) classical definition of Probability:-

An experiment may results n exhaustive mutual exclusive. In which of them " m " are favourable to an event " E " then Probability of getting an event E is

$$P(E) = \frac{\text{Favourable Cases}}{\text{Total exhaustive Cases}} = \frac{m}{n}$$

$$P(E) = \frac{m}{n}$$

Total Probability = 1

$$P(E) + P(\bar{E}) = 1$$

$$P(\bar{E}) = 1 - P(E)$$

$$P(\bar{E}) = 1 - \frac{m}{n}$$

Limits of Probability

$$0 \leq P \leq 1$$

(i) $P(E) = 1$ then that event is called Certain event!

(ii) $P(E) = 0$ then that event is called impossible event-



Problems -

① Two dice are rolled. what is the Probability of getting Sum is 9.

Sol:- Favourable Cases for getting Sum 9 is
 $\{(5,4) (4,5) (3,6) (6,3)\} = 4$

Total Exhaustive Case for two dice are rolled = $6^2 = 36$.

Prob of getting Sum is 9 = $\frac{\text{Favourable cases}}{\text{Total exhaustive cases}} = \frac{4}{36}$.

② A card is drawn from well shuffled pack what is the Prob of getting a King.

Sol:- Favourable Cases for getting a King = $4C_1$.

Total exhaustive Cases for drawing a card = $52C_1$.

Prob of getting a King = $\frac{\text{Favourable cases}}{\text{Total exhaustive case}} = \frac{4C_1}{52C_1}$

$$= \frac{4}{52}$$

③ Two Coins are rolled what is the Prob of getting (i) Both heads (ii) both tails (iii) one head.

Sol:- Two Coins are rolled Total exhaustive Cases = $2^2 = 4$.

$\{(HH) (TT) (HT) (TH)\}$.

(i) Favourable Cases for both heads = 1

Prob of getting both heads = $\frac{F.C.}{T.E.C.} = \frac{1}{4}$



(ii) Favourable Cases for both tails = 1

$$\text{Prob of getting both tails} = \frac{F.C.}{T.E.C} = \frac{1}{4}$$

(iii) Favourable Cases for one head = 2

$$\text{Prob of getting one head} = \frac{F.C.}{T.E.C} = \frac{2}{4}$$

Problem (4): A stockist has 20 items in a lot of which 12 are non defectives, if customer selects 3 items from a lot.

(i) What is the Prob of 3 are Non defectives.

(ii) What is the Prob of 2 are non defective and one is defective.

Sol:- (i) Total exhaustive Cases for drawing three items = ${}^{20}C_3$

Favourable Cases for drawing 3 Non defectives = ${}^{12}C_3$

$$\text{Prob of getting 3 are non defectives} = \frac{F.C.}{T.E.C} = \frac{{}^{12}C_3}{{}^{20}C_3}$$

(ii) favourable cases for two are non defective and one is defective =

$${}^{12}C_2 \cdot {}^8C_1$$

Total Exhaustive Cases = ${}^{20}C_3$



Prob of getting two are non defective and one is defective = $\frac{F \cdot C}{T \cdot E \cdot C} = \frac{{}^8P_2 \cdot {}^8C_1}{{}^{20}C_3}$

→ Conditional Probability:-

If A and B are two dependent events then,

$$P(A/B) = \frac{P(A \cap B)}{P(B)}$$

ie Prob of occurrence of event A, when event B is already happened.

$$P(B/A) = \frac{P(A \cap B)}{P(A)}$$

ie Prob of occurrence of event B, when event A is already happened.

Note:- If $A_1, A_2, A_3, \dots, A_n$ are Independent events the prob of getting at least one of the event is

$$\begin{aligned} P(A_1 \cup A_2 \cup A_3 \dots \cup A_n) &= 1 - P(\overline{A_1 \cup A_2 \cup A_3 \dots \cup A_n}) \\ &= 1 - P(\overline{A_1} \cap \overline{A_2} \cap \overline{A_3} \cap \dots \cap \overline{A_n}) \\ &= 1 - [P(\overline{A_1}) \cdot P(\overline{A_2}) \cdot P(\overline{A_3}) \dots P(\overline{A_n})] \end{aligned}$$



Problem :- (1) A problem in statistics given to three students A, B and C whose chances are solving are $\frac{1}{2}$, $\frac{1}{4}$ and $\frac{1}{4}$. What is the prob of the problem will be solve atleast one of them, ~~are~~ All three students try Independently.

Sol:- Prob of that problem solved by student A $P(A) = \frac{1}{2}$

Prob of that problem solved by student B $P(B) = \frac{1}{4}$.

Prob of that problem solved by student C $P(C) = \frac{1}{4}$.

$$P(\bar{A}) = 1 - P(A) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$P(\bar{B}) = 1 - P(B) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$P(\bar{C}) = 1 - P(C) = 1 - \frac{1}{4} = \frac{3}{4}$$

Prob of that problem solved atleast by one of them is

$$P(A \cup B \cup C) = 1 - [P(\bar{A}) \cdot P(\bar{B}) \cdot P(\bar{C})]$$

$$= 1 - \left(\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{3}{4} \right)$$

$$= 1 - \frac{9}{32} = \frac{23}{32}$$

Problem :- (2) The prob of A will live up to 60 years is $\frac{3}{4}$. Prob of B will live up to 60 years is $\frac{2}{3}$. What is the prob that

- (i) Both A and B are live up to 60 years.
- (ii) Both die before reaching 60 years.



Sol: Prob of A will live up to 60 yrs $P(A) = \frac{3}{4}$.

Prob of B will live up to 60 yrs $P(B) = \frac{2}{3}$.

(i) Both A and B are live up to 60 years

$$P(A \cap B) = P(A) \cdot P(B)$$

$$= \frac{3}{4} \cdot \frac{2}{3} = \frac{6}{12} \Rightarrow \frac{1}{2}$$

(ii) Both die before reaching 60 years $P(A \cup B) = 1 - P(\overline{A \cup B})$

$$= 1 - \{P(\overline{A}) \cdot P(\overline{B})\}$$

$$P(\overline{A}) = 1 - P(A)$$

$$= 1 - \frac{3}{4} = \frac{1}{4}$$

$$P(\overline{B}) = 1 - P(B)$$

$$= 1 - \frac{2}{3} = \frac{1}{3}$$

$$P(A \cup B) = 1 - \{P(\overline{A}) \cdot P(\overline{B})\}$$

$$= 1 - \left\{ \frac{1}{4} \cdot \frac{1}{3} \right\}$$

$$= 1 - \frac{1}{12}$$

$$= \frac{11}{12}$$

Problem 3 A Bag contains 25 tickets numbered 1 to 25. one ticket is drawn what is the prob of getting multiple of 2 or 5.

Sol: let A_1 be the multiple of 2 and A_2 be the multiple of 5.

Total exhaustive cases for drawing a ticket

$$n = 25$$



Favourable Cases for multiple of 2 is

$$= \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24\} = 12_c$$

Prob of getting a ticket and Multiple of 2 is

$$P(A_1) = \frac{12_c}{25_c} = \frac{12}{25}$$

Favourable Cases for Multiple of 5 is

$$\{5, 10, 15, 20, 25\} = 5_c$$

Prob of getting a ticket and Multiple of 5 is

$$P(A_2) = \frac{5_c}{25_c} = \frac{5}{25}$$

Favourable Cases for Multiple of 2 and 5 is

$$\{10, 20\} = 2_c$$

Prob of getting Multiple of 2 and 5 are

$$P(A_1 \cap A_2) = \frac{2_c}{25_c} = \frac{2}{25}$$

Prob of getting Multiple of 2 (or) 5 is

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$$

$$= \frac{12}{25} + \frac{5}{25} - \frac{2}{25}$$

$$P(A_1 \cup A_2) = \frac{15}{25} = \frac{3}{5}$$

$$= 0.6$$



Theorem

Addition theorem on Probability for $n=2$ events

Statement:- If A_1 and A_2 are two disjoint events. then.

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2).$$

Proof:- From Venn diagram.

$$A_1 \cup A_2 = A_1 \cup (A_2 \cap \bar{A}_1)$$

Taking probabilities on both sides

$$P(A_1 \cup A_2) = P(A_1 \cup (A_2 \cap \bar{A}_1)).$$

$$P(A_1 \cup A_2) = P(A_1) + P(A_2 \cap \bar{A}_1) \quad \text{--- (1)} \quad [\because A_1, A_2 \text{ are disjoint sets }].$$

$$\text{also } A_2 = (A_1 \cap A_2) \cup (A_2 \cap \bar{A}_1)$$

Taking probabilities on both sides.

$$P(A_2) = P[(A_1 \cap A_2) \cup (A_2 \cap \bar{A}_1)].$$

$$P(A_2) = P(A_1 \cap A_2) + P(A_2 \cap \bar{A}_1) \quad \text{--- (2)}$$

$$P(A_2) - P(A_1 \cap A_2) = P(A_2 \cap \bar{A}_1) \quad \text{--- (2)}$$

Substitute equation (2) in to equ (1).

$$P(A_1 \cup A_2) = P(A_1) + P(A_2 \cap \bar{A}_1)$$

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$$

Hence the proof.



Theory Addition Theorem on Probability for n-Events:-

Statement:- $A_1, A_2, A_3, \dots, A_n$ are n events has been drawn from sample space. Then.

$$P\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) + (-1)^{n-1} P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n).$$

(or)

$$P(A_1 \cup A_2 \cup A_3 \dots \cup A_n) = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) - \dots + (-1)^{n-1} P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n)$$

Proof:- This can be proved it by mathematical induction.

For $n=2$ events.

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$$

It is true for $n=2$ events.

For $n=3$ events.

$$P(\underbrace{A_1}_{A_1} \cup \underbrace{A_2 \cup A_3}_{A_2}) = P(A_1) + P(A_2 \cup A_3) - P\left[\underbrace{A_1 \cap (A_2 \cup A_3)}_{A_1 \cap A_2 \cup A_1 \cap A_3}\right]$$

$$= P(A_1) + P(A_2 \cup A_3) - P\left[\underbrace{(A_1 \cap A_2)}_{A_1} \cup \underbrace{(A_1 \cap A_3)}_{A_2}\right]$$

$$= P(A_1) + P(A_2 \cup A_3) - \left[P(A_1 \cap A_2) + P(A_1 \cap A_3) - P(A_1 \cap A_2 \cap A_3) \right]$$



$$= P(A_1) + P(A_2 \cup A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) + P(A_1 \cap A_2 \cap A_3)$$

$$= P(A_1) + P(A_2) + P(A_3) - P(A_2 \cap A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) + P(A_1 \cap A_2 \cap A_3)$$

$$= P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_2 \cap A_3) - P(A_1 \cap A_3) + P(A_1 \cap A_2 \cap A_3)$$

It is true for $n=3$ events.

Let us assume that the given statement is true for $n=r$ events.

$$P\left[\bigcup_{i=1}^n A_i\right] = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) - \dots$$

$$(-1)^{r-1} \cdot P(A_1 \cap A_2 \cap A_3 \dots A_r)$$

For $n=r+1$ events

$$P\left(\bigcup_{i=1}^{r+1} A_i\right) = P\left[\underbrace{\bigcup_{i=1}^r A_i}_{A_1} \cup \underbrace{A_{r+1}}_{A_2}\right]$$

$$= P\left[\bigcup_{i=1}^r A_i\right] + P(A_{r+1}) - P\left[\bigcup_{i=1}^r A_i \cap A_{r+1}\right]$$

$$= \sum_{i=1}^r P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) - \dots$$

$$(-1)^{r-1} \cdot P(A_1 \cap A_2 \cap \dots \cap A_r) + P(A_{r+1}) - P\left(\bigcup_{i=1}^r A_i \cap A_{r+1}\right)$$



$$= \sum_{i=1}^{n+1} P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) \dots$$

$$(-1)^{n+1} \cdot P(A_1 \cap A_2 \cap A_3 \dots A_n) - \left[\sum_{i=1}^n P(A_i \cap A_{n+1}) - \sum_{i < j} P(A_i \cap A_j \cap A_{n+1}) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k \cap A_{n+1}) \dots \right]$$

$$\sum_{i < j} P(A_i \cap A_j \cap A_{n+1}) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k \cap A_{n+1}) \dots$$

$$(-1)^{n+1} \cdot P(A_1 \cap A_2 \cap A_3 \cap \dots A_n \cap A_{n+1})$$

It is true for n = 2 events.

It is also true for n = 1 events. According to the Principle of Mathematical Induction the given statement is true for all positive integers.

Hence the proof.

Theorem

Multiplication theorem on Probability

for n-events:-

Statement:- $A_1, A_2, A_3, \dots, A_n$ are n events defined on a sample space then

$$P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2/A_1) \cdot P(A_3/A_1 \cap A_2) \dots \dots \dots P(A_n/A_1 \cap A_2 \cap A_3 \dots \dots A_{n-1})$$



Proof:- This can be proved it by mathematical induction.

For $n=2$ events

according to Conditional Probability

$$P(A_1 \cap A_2) = P(A_1) \cdot P(A_2|A_1)$$

It is true for $n=2$ events.

For $n=3$ events

$$P(A_1 \cap A_2 \cap A_3) = P(A_1 \cap A_2) \cdot P(A_3|A_1 \cap A_2)$$

$$= P(A_1) \cdot P(A_2|A_1) \cdot P(A_3|A_1 \cap A_2)$$

It is true for $n=3$ events.

Let us take the given statement is true for $n=r$ events

$$P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_r) = P(A_1) \cdot P(A_2|A_1) \cdot P(A_3|A_1 \cap A_2) \cdot \dots \cdot P(A_r|A_1 \cap A_2 \cap A_3 \cap \dots \cap A_{r-1})$$

For $n=r+1$ events

$$P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_{r+1}) = P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_r \cap A_{r+1})$$

$$= P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_r) \cdot P(A_{r+1}|A_1 \cap A_2 \cap A_3 \cap \dots \cap A_r)$$

$$P(A_1) \cdot P(A_2|A_1) \cdot P(A_3|A_1 \cap A_2) \cdot \dots \cdot P(A_r|A_1 \cap A_2 \cap \dots \cap A_{r-1}) \cdot P(A_{r+1}|A_1 \cap A_2 \cap \dots \cap A_r)$$



$$= P(A_1) \cdot P(A_2/A_1) \cdot P(A_3/A_1 A_2) \cdots P(A_n/A_1 A_2 \cdots A_{n-1})$$

It is true for $n = r+1$ events


It is also true for $n = r$ events

According to Mathematical Induction
the given statement is true for all
Positive Integers.

Hence the proof



RANDOM VARIABLES



Random Variable :- A real valued function defined on a sample space is called Random Variable. It can take the values on real line $(-\infty \text{ to } +\infty)$

(or)
Random Variable means a real number x can be associated with each outcome of a random experiment. It can take any one of the several possible outcomes with definite probability.

EX!- (i) In tossing a coin getting a head is denoted with random variable x , it can take the values $x = \{0, 1\}$.

its sample space $S = \{H, T\}$.

random variable x is getting head $x = \{0, 1\}$.

EX!- In tossing two coins getting a head is denoted with random variable x .



It can take the values $X = \{0, 1, 2\}$.
 A sample space $S = \{HH, TT, TH, HT\}$.
 Random variable X is getting head $X = \{0, 1, 2\}$.

Discrete Random Variable:— A random variable X can assume at most countable number of values (or) finite number of values. then it is called discrete random variable. (or)

A real valued function defined on a discrete sample space is called discrete random variable.

Continuous Random Variable:— A random variable X can assume infinite (or) uncountable number of values then it is called continuous random variable. (or)

A real valued function defined on a continuous sample space is called continuous random variable.

Probability mass function (P.M.F):—
 Suppose X is discrete random variable taking at most countable



number of values i.e. $x_1, x_2, x_3, \dots, x_n$
 if some Probability $P(x_1), P(x_2), P(x_3), \dots, P(x_n)$
 are must satisfies the following conditions.

(i) $P(x_i) \geq 0$

(ii) $\sum_{i=1}^n P(x_i) = 1$

The function $P(x_i)$ is called Probability mass
 function of the random variable X .

x_i	x_1	x_2	x_3	\dots	x_n
$P(x_i)$	$P(x_1)$	$P(x_2)$	$P(x_3)$	\dots	$P(x_n)$

Mean $\mu_1 = \sum_{i=1}^n x_i P(x_i)$

$\mu_2 = \sum_{i=1}^n x_i^2 P(x_i)$

Variance $\mu_2 = \mu_2' - (\mu_1')^2$

Variance $\mu_2 = \sum_{i=1}^n x_i^2 P(x_i) - \left[\sum_{i=1}^n x_i P(x_i) \right]^2$

Probability density function (P.d.f):-

consider the small interval
 $[x - \frac{1}{2}dx, x + \frac{1}{2}dx]$ of length of dx around the
 point x . $f(x)$ is any continuous function of x
 such that $f(x) dx$ represented the Probability
 density function of random variable x . It
 must satisfies the following conditions.

(1) $f(x) \geq 0$

(2) $\int_{-\infty}^{\infty} f(x) dx = 1$



$$\text{Mean } \mu_1 = \int_{-a}^a x \cdot f(x) dx$$

$$\mu_2 = \int_{-a}^a x^2 f(x) dx$$

$$\begin{aligned} \text{Variance } \mu_2 &= \mu_2 - (\mu_1)^2 \\ &= \int_{-a}^a x^2 f(x) dx - \left[\int_{-a}^a x f(x) dx \right]^2 \end{aligned}$$

Problems

- ① A random Variable x has the following Probability function.

$$x: 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \quad 7 \dots$$

$$P(x): 0 \quad k \quad 2k \quad 2k \quad 3k \quad k^2 \quad 2k^2 \quad 7k^2 + k \dots$$

- (i) Find the Value of k . (ii) $P(x < 6)$ (iii) $P(x > 6)$.
(iv) $P(0 < x < 5)$.

Sol:- We know that $\sum P(x_i) = 1$

$$0 + k + 2k + 2k + 3k + k^2 + 2k^2 + 7k^2 + k = 1$$

$$10k^2 + 9k = 1$$

$$10k^2 + 9k - 1 = 0$$

$$10k^2 + 10k - k - 1 = 0$$

$$10k[k+1] - 1[k+1] = 0$$

$$(k+1) = 0 \quad \left| \quad \begin{array}{l} 10k - 1 = 0 \\ 10k = 1 \end{array} \right.$$

$$\boxed{k = -1}$$

$$k = \frac{1}{10}$$

$$\therefore k = \frac{1}{10}$$

$x:$	0	1	2	3	4	5	6	7
$P(x):$	0	0.1	0.2	0.2	0.3	0.01	0.02	0.17

$$\begin{aligned} \text{(ii)} \quad P[X < 6] &= P[X=0] + P[X=1] + P[X=2] + P[X=3] + P[X=4] + P[X=5] \\ &= 0 + 0.1 + 0.2 + 0.2 + 0.3 + 0.01 \\ &= 0.81 \checkmark \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad P[X > 6] &= P[X=6] + P[X=7] \\ &= 0.02 + 0.17 \\ &= 0.19 \end{aligned}$$

$$\begin{aligned} P[X > 6] &= 1 - P[X < 6] \\ &= 1 - 0.81 \\ &= 0.19 \end{aligned}$$

$$\begin{aligned} P[X > a] + P[X < a] &= 1 \\ P[X > a] &= 1 - P[X < a] \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad P[0 < X < 5] &= P[X=1] + P[X=2] + P[X=3] + P[X=4] \\ &= 0.1 + 0.2 + 0.2 + 0.3 \\ &= \underline{\underline{0.8}} \end{aligned}$$

Problem 2 In throwing a dice find probability distribution and also find

$$\text{(i)} P[X > 5] \quad \text{(ii)} P[X < 5] \quad \text{(iii)} P[X > 2] \quad \text{(iv)} P[0 \leq X \leq 4]$$

Sol A dice was thrown Prob distribution becomes.

$x:$	1	2	3	4	5	6
$P(x):$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

$$\begin{aligned} \text{(i)} \quad P[X > 5] &= P[X=5] + P[X=6] \\ &= \frac{1}{6} + \frac{1}{6} = \frac{2}{6} = \frac{1}{3} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad P[X < 5] &= 1 - P[X > 5] \\ &= 1 - \frac{1}{3} = \frac{2}{3} \end{aligned}$$



$$(ii) P[X > 2] = P[X=2] + P[X=3] + P[X=4] + P[X=5] + P[X=6]$$

$$= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6}$$

$$= \frac{5}{6}$$

$$(iv) P[0 \leq X \leq 4] = P[X=1] + P[X=2] + P[X=3] + P[X=4]$$

$$= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6}$$

$$= \frac{4}{6} = \frac{2}{3}$$

Problem

- (3) Two dice are thrown find the Prob. distribution of sum of the faces and also find (i) $P[X > 4]$ (ii) $P[X \leq 6]$ (iii) $P[1 \leq X < 6]$ and also find Mean and Variance.

Sol:- Probability distribution of Sum of two dice is

Sum	Favourable Cases	Prob.
2	(1,1)	$\frac{1}{36}$
3	(1,2) (2,1)	$\frac{2}{36}$
4	(3,1) (1,3) (2,2)	$\frac{3}{36}$
5	(2,3) (3,2) (4,1) (1,4)	$\frac{4}{36}$
6	(3,3) (5,1) (1,5) (4,2) (2,4)	$\frac{5}{36}$
7	(1,6) (6,1) (5,2) (2,5) (4,3) (3,4)	$\frac{6}{36}$
8	(6,2) (2,6) (4,4) (5,3) (3,5)	$\frac{5}{36}$
9	(5,4) (4,5) (6,3) (3,6)	$\frac{4}{36}$
10	(5,5) (6,4) (4,6)	$\frac{3}{36}$
11	(6,5) (5,6)	$\frac{2}{36}$
12	(6,6)	$\frac{1}{36}$



Prob distribution

$x:$	2	3	4	5	6	7	8	9	10	11	12
$P(x)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

$$\begin{aligned} \text{(i)} \quad P[x > 4] &= P(x=4) + P(x=5) + P(x=6) + P(x=7) + P(x=8) + P(x=9) + P(x=10) \\ &\quad + P(x=11) + P(x=12) \\ &= \frac{3}{36} + \frac{4}{36} + \frac{5}{36} + \frac{6}{36} + \frac{5}{36} + \frac{4}{36} + \frac{3}{36} + \frac{2}{36} \\ &= \frac{33}{36} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad P[x > 4] &= 1 - P[x < 4] \\ &= 1 - \{P(x=2) + P(x=3)\} \\ &= 1 - \left\{ \frac{1}{36} + \frac{2}{36} \right\} \\ &= 1 - \frac{3}{36} \\ &= \frac{33}{36} \end{aligned}$$

$$P[x > 4] = \frac{31}{36}$$

$$\begin{aligned} \text{(iii)} \quad P[x \leq 6] &= P(x=2) + P(x=3) + P(x=4) + P(x=5) + P(x=6) \\ &= \frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{4}{36} + \frac{5}{36} \\ &= \frac{15}{36} \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad P[1 \leq x < 6] &= P(x=2) + P(x=3) + P(x=4) + P(x=5) \\ &= \frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{4}{36} \\ &= \frac{10}{36} \end{aligned}$$

$$\text{(iv) Mean} \quad \mu_1 = \sum x \cdot P(x)$$

$$\begin{aligned} \text{Mean}(\mu_1) &= (2) \left(\frac{1}{36}\right) + (3) \left(\frac{2}{36}\right) + (4) \left(\frac{3}{36}\right) + (5) \left(\frac{4}{36}\right) + (6) \left(\frac{5}{36}\right) + (7) \left(\frac{6}{36}\right) \\ &\quad + (8) \left(\frac{5}{36}\right) + (9) \left(\frac{4}{36}\right) + (10) \left(\frac{3}{36}\right) + (11) \left(\frac{2}{36}\right) + (12) \left(\frac{1}{36}\right) \\ &= \frac{2}{36} + \frac{6}{36} + \frac{12}{36} + \frac{20}{36} + \frac{30}{36} + \frac{42}{36} + \frac{40}{36} + \frac{36}{36} + \frac{30}{36} + \frac{22}{36} + \frac{12}{36} \end{aligned}$$

$$\text{Mean}(\mu_1) = \frac{252}{36} = 7$$



$$\therefore \text{mean } \mu_1 = 7$$

$$\text{Variance } \mu_2 = \mu_3' - (\mu_1')^2$$

$$\mu_3' = \sum x^2 P(x)$$

$$= (2)^2 \left(\frac{1}{36}\right) + (3)^2 \left(\frac{2}{36}\right) + (4)^2 \left(\frac{3}{36}\right) + (5)^2 \left(\frac{4}{36}\right) + (6)^2 \left(\frac{5}{36}\right) + (7)^2 \left(\frac{6}{36}\right) \\ + (8)^2 \left(\frac{5}{36}\right) + (9)^2 \left(\frac{4}{36}\right) + (10)^2 \left(\frac{3}{36}\right) + (11)^2 \left(\frac{2}{36}\right) + (12)^2 \left(\frac{1}{36}\right)$$

$$= \frac{4}{36} + \frac{18}{36} + \frac{48}{36} + \frac{100}{36} + \frac{180}{36} + \frac{294}{36} + \frac{320}{36} + \frac{324}{36} + \frac{300}{36} \\ + \frac{242}{36} + \frac{144}{36}$$

$$\mu_3' = \frac{1974}{36}$$

$$\mu_3' = 54.8333$$

$$\text{Variance} = \mu_3' - (\mu_1')^2$$

$$= 54.8333 - (7)^2$$

$$= 54.8333 - 49$$

$$\text{Variance } \mu_2 = 5.8333 \quad \checkmark$$



Problem (3) If the Random Variable X has the following Prob mass function.

$$P[X=r] = P(r) = \frac{1}{2^r}, \quad r=1, 2, 3, 4, \dots$$

find (i) $P[r \text{ is even}]$ (ii) $P[r > 5]$.

Sol - Given Prob mass function $P[X=r] = P(r) = \frac{1}{2^r}$

$$r=1, 2, 3, 4, \dots$$

$$P(r) = \frac{1}{2^1}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \dots$$

$$\begin{aligned} \text{(i) } P[r \text{ is even}] &= P[r=2] + P[r=4] + P[r=6] + \dots \\ &= \frac{1}{2^2} + \frac{1}{2^4} + \frac{1}{2^6} + \dots \end{aligned}$$

Sum of Infinite Series

'a' is first term
'r' is common ratio.

$$S_{\infty} = \frac{a}{1-r}$$

$$\text{Here } a = \frac{1}{2}, \quad r = \frac{r_2}{r_1} = \frac{\frac{1}{2^4}}{\frac{1}{2^2}} = \frac{1}{2^4} \times \frac{2^2}{1}$$

$$r = \frac{1}{2^2} = \frac{1}{4}$$

$$S_{\infty} = \frac{a}{1-r} = \frac{\frac{1}{2}}{1-\frac{1}{4}} = \frac{\frac{1}{2}}{\frac{3}{4}} = \frac{1}{2} \times \frac{4}{3} = \frac{2}{3}$$

$$\text{(ii) } P[r > 5] = 1 - P[r < 5]$$

$$= 1 - \{P(r=1) + P(r=2) + P(r=3) + P(r=4)\}$$

$$= 1 - \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16}$$

$$= 1 - 0.9375$$

$$= 0.0625$$

$$P[X > 5] = 0.0625$$



Problem 4 Probability mass function of random variable x is

$$P(x=x) = P(x) = \frac{x}{15}, \quad x=1, 2, 3, 4 \text{ and } 5.$$

Find (i) $P(x=1 \text{ (or) } x=2)$ (ii) $P\left[\frac{1}{2} < x < \frac{5}{2} \mid x > 1\right]$.

(iii) $P\left[\frac{1}{2} < x < \frac{5}{2}\right]$.

Sol Given Prob mass function of random variable x is

$$P(x=x) = P(x) = \frac{x}{15}, \quad x=1, 2, 3, 4 \text{ and } 5.$$

$$x: \quad 1 \quad 2 \quad 3 \quad 4 \quad 5$$

$$P(x): \quad \frac{1}{15} \quad \frac{2}{15} \quad \frac{3}{15} \quad \frac{4}{15} \quad \frac{5}{15}$$

(i) $P(x=1 \text{ (or) } x=2) = P(x=1) + P(x=2)$

$$= \frac{1}{15} + \frac{2}{15}$$

$$= \frac{3}{15} = \frac{1}{5}$$

(ii) $P\left[\frac{1}{2} < x < \frac{5}{2} \mid x > 1\right] = P[0.5 < x < 2.5 \mid x > 1]$

x lies between 0.5 and 2.5 but it is greater than 1.

$$\therefore P(x=2) = \frac{2}{15}$$

(iii) $P\left[\frac{1}{2} < x < \frac{5}{2}\right] = P[0.5 < x < 2.5]$

$$= P(x=1) + P(x=2)$$

$$= \frac{1}{15} + \frac{2}{15} = \frac{3}{15}$$

$$= \frac{1}{5}$$



Problem:- In rolling a dice the number of faces is denoted with random variable x . Then write the probability distribution. also find Mean and Variance.

Sol. - A dice is rolled Prob distribution becomes:

x_i :	1	2	3	4	5	6
$P(x_i)$:	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

$$\text{Mean } \mu_1 = \sum x_i P(x_i)$$

$$= (1)\left(\frac{1}{6}\right) + (2)\left(\frac{1}{6}\right) + (3)\left(\frac{1}{6}\right) + (4)\left(\frac{1}{6}\right) + (5)\left(\frac{1}{6}\right) + (6)\left(\frac{1}{6}\right)$$

$$= \frac{21}{6}$$

$$\text{Q.2 - Variance } \mu_2 = \mu_2' - (\mu_1')^2$$

$$\mu_2' = \sum x_i^2 P(x_i)$$

$$= (1)^2\left(\frac{1}{6}\right) + (2)^2\left(\frac{1}{6}\right) + (3)^2\left(\frac{1}{6}\right) + (4)^2\left(\frac{1}{6}\right) + (5)^2\left(\frac{1}{6}\right) + (6)^2\left(\frac{1}{6}\right)$$

$$= \frac{1}{6} + \frac{4}{6} + \frac{9}{6} + \frac{16}{6} + \frac{25}{6} + \frac{36}{6}$$

$$\mu_2' = \frac{91}{6}$$

$$\text{Variance } \mu_2 = \mu_2' - (\mu_1')^2$$

$$= \frac{91}{6} - \left(\frac{21}{6}\right)^2$$

$$= \frac{91}{6} - \frac{441}{36}$$

$$\text{Variance } \mu_2 = \frac{546 - 441}{36} = \frac{105}{36}$$



Note:- If x is a continuous random variable. $f(x)$ is a prob density function lies between a to b . then

① n^{th} central moment $\mu_n = \int_a^b (x-\bar{x})^n \cdot f(x) dx$.

② n^{th} raw moment about the point A is

$$\mu_n' = \int_a^b (x-A)^n \cdot f(x) dx$$

③ n^{th} raw moment about origin,

$$\mu_n' = \int_a^b (x-0)^n \cdot f(x) dx$$

$$\mu_n' = \int_a^b x^n \cdot f(x) dx \quad \text{--- ①}$$

Put $n=1$ in eqn ① $\mu_1' = \int_a^b x^1 \cdot f(x) dx = \text{Mean}$.

Put $n=2$ in eqn ① $\mu_2' = \int_a^b x^2 \cdot f(x) dx$.

~~Put~~ Variance $\mu_2 = \mu_2' - (\mu_1')^2$

$$\mu_2 = \int_a^b x^2 \cdot f(x) dx - \left[\int_a^b x \cdot f(x) dx \right]^2$$

Put $n=3$ in eqn ① $\mu_3' = \int_a^b x^3 \cdot f(x) dx$.

Put $n=4$ in eqn ① $\mu_4' = \int_a^b x^4 \cdot f(x) dx$.

Problems:

① If x is a continuous random variable with prob density function $f(x) = Ax^2, 0 \leq x \leq 1$
~~obtain~~ Determine the value of A and find mean and variance.

Sol. - Given prob density function $f(x) = Ax^2, 0 \leq x \leq 1$

We know that $\int_a^b f(x) dx = 1$

$$\int_0^1 Ax^2 dx = 1$$

$$A \int_0^1 x^2 dx = 1$$

$$A \left[\frac{x^3}{3} \right]_0^1 = 1$$

$$\frac{A}{3} [1 - 0] = 1$$

$$\boxed{A = 3}$$

\therefore Prob density function $f(x) = 3x^2, 0 \leq x \leq 1$

$$\text{Mean } \mu_1' = \int_0^1 x \cdot f(x) dx$$

$$= \int_0^1 x \cdot (3x^2) dx$$

$$= 3 \int_0^1 x^3 dx$$

$$= 3 \left[\frac{x^4}{4} \right]_0^1$$

$$= \frac{3}{4} [x^4]_0^1$$

$$\text{Mean } \mu_1' = \frac{3}{4} [1 - 0] = \frac{3}{4}$$

$$\therefore \text{Mean } \mu_1 = 3/4$$

$$\text{Variance } \mu_2 = \mu_2' - (\mu_1')^2$$

$$\begin{aligned} \mu_2' &= \int_a^b x^2 \cdot f(x) dx \\ &= \int_0^1 x^2 \cdot (3x^2) dx \\ &= 3 \int_0^1 x^4 dx \\ &= 3 \left[\frac{x^5}{5} \right]_0^1 \end{aligned}$$

$$\mu_2' = \frac{3}{5} [1-0] = 3/5$$

$$\begin{aligned} \therefore \text{Variance } \mu_2 &= \mu_2' - (\mu_1')^2 \\ &= \frac{3}{5} - \left(\frac{3}{4}\right)^2 \\ &= \frac{3}{5} - \frac{9}{16} \\ &= \frac{48-45}{80} = 0.375 \end{aligned}$$

$$\text{Variance } \mu_2 = 0.375$$

Problem (2) The diameter of the electric cable say x is assumed to be a continuous random variable with prob density function $f(x) = 6x(1-x)$, $0 \leq x \leq 1$, check that cable is prob density function also calculate mean and variance.

Sol Given prob density function $f(x) = 6x(1-x)$
 $0 \leq x \leq 1$



To check given function is prob density function or not we have to prove total prob = 1

$$\begin{aligned}\int_0^1 f(x) dx &= \int_0^1 6x(1-x) dx \\ &= 6 \int_0^1 (x - x^2) dx \\ &= 6 \int_0^1 x \cdot dx - \int_0^1 x^2 \cdot dx \\ &= 6 \left[\frac{x^2}{2} \right]_0^1 - \left[\frac{x^3}{3} \right]_0^1 \\ &= \frac{6}{2} [1-0] - \frac{6}{3} [1-0] \\ &= \frac{6^3}{2} - \frac{6^2}{3} \\ &= 3-2\end{aligned}$$

$$\int_0^1 f(x) dx = 1$$

\therefore Given function is Prob density

function.

$$\therefore \text{mean } \mu_1 = \int_0^1 x \cdot f(x) dx$$

$$\begin{aligned}\mu_1 &= \int_0^1 x \cdot 6x(1-x) dx \\ &= 6 \int_0^1 x^2(1-x) dx \\ &= 6 \left[\int_0^1 x^2 \cdot dx - \int_0^1 x^3 \cdot dx \right] \\ &= 6 \left[\left[\frac{x^3}{3} \right]_0^1 - \left[\frac{x^4}{4} \right]_0^1 \right] \\ &= 6 \left[\left(\frac{1}{3} - 0 \right) - \left(\frac{1}{4} - 0 \right) \right] \\ &= 6 \left[\frac{4-3}{12} \right] = \frac{1}{2}\end{aligned}$$



$$\therefore \text{Mean } \mu_1 = \frac{1}{2}$$

$$\text{Variance } \mu_2 = \mu_2' - (\mu_1)^2$$

$$\mu_2' = \int_0^1 x^2 \cdot f(x) dx$$

$$= \int_0^1 x^2 \cdot 6x(1-x) dx$$

$$= 6 \left[\int_0^1 x^3(1-x) dx \right]$$

$$= 6 \left[\int_0^1 (x^3 - x^4) dx \right]$$

$$= 6 \left[\int_0^1 x^3 dx - \int_0^1 x^4 dx \right]$$

$$= 6 \left[\left[\frac{x^4}{4} \right]_0^1 - \left[\frac{x^5}{5} \right]_0^1 \right]$$

$$= 6 \left[\left(\frac{1}{4} - 0 \right) - \left(\frac{1}{5} - 0 \right) \right]$$

$$= 6 \left[\frac{1}{4} - \frac{1}{5} \right]$$

$$= 6 \left[\frac{5-4}{20} \right]$$

$$\mu_2' = \frac{6}{20}$$

$$\therefore \text{Variance } \mu_2 = \mu_2' - (\mu_1)^2$$

$$= \frac{6}{20} - \left(\frac{1}{2} \right)^2$$

$$= \frac{6}{20} - \frac{1}{4}$$

$$\text{Variance} = \frac{6-5}{20} = \frac{1}{20} = 0.05$$

$$\frac{4}{11.5}$$



Problem 3 If x is continuous random variable with prob density function

$$f(x) = \begin{cases} ax & 0 \leq x \leq 1 \\ a & 1 \leq x \leq 2 \\ -ax + 3a & 2 \leq x \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

(i) Find the constant value "a"

(ii) $P[x \leq 1.5]$

Sol:- (i) To find constant "a"

We know that $\int_{-\infty}^{\infty} f(x) dx = 1$

$$\int_{-\infty}^0 f(x) dx + \int_0^1 f(x) dx + \int_1^2 f(x) dx + \int_2^3 f(x) dx + \int_3^{\infty} f(x) dx = 1$$

$$0 + \int_0^1 ax \cdot dx + \int_1^2 a \cdot dx + \int_2^3 (-ax + 3a) dx + 0 = 1$$

$$= a \left[\frac{x^2}{2} \right]_0^1 + a [x]_1^2 - a \left[\frac{x^2}{2} \right]_2^3 + 3a [x]_2^3 = 1$$

$$= \frac{a}{2} [1-0] + a[2-1] - \frac{a}{2} [9-4] + 3a[3-2] = 1$$

$$= \frac{a}{2} + a - \frac{5a}{2} + 3a = 1$$

$$= \cancel{a} + \cancel{a} - \frac{5a}{2} + \cancel{3a} = 1$$

$$\frac{a + 2a - 5a + 6a}{2} = 1$$

$$= \frac{4a}{2} = 1$$

$$= \frac{4a}{2} = 1$$

$$2a = 1$$

$$\boxed{a = \frac{1}{2}}$$



$$(ii) P[X \leq 1.5].$$

$$= \int_{-\infty}^0 f(x) dx + \int_0^1 f(x) dx + \int_1^{1.5} f(x) dx.$$

$$= 0 + \int_0^1 ax \cdot dx + \int_1^{1.5} a \cdot dx.$$

$$= 0 + a \left[\frac{x^2}{2} \right]_0^1 + a [x]_1^{1.5}.$$

$$= \frac{a}{2} [1-0] + a [1.5-1].$$

$$= \frac{a}{2} + 0.5a.$$

$$P[X \leq 1] = \frac{a}{2} + \frac{a}{2} = \frac{2a}{2}$$

$$P[X \leq 1.5] = \frac{1}{2}$$

Problem,

→ If x is a continuous r.v. with

prob density function $f(x) = 6x^2 e(1-x)$, $0 \leq x \leq 1$

find constant " e ". also find mean

and variance.

<p style="text-align: center;"><u>Ans</u></p> <p>$e = 2$</p> <p>mean $\mu_1 = 3/5$</p> <p>$\mu_2 = 2/5$</p> <p>Variance $\mu_2 = 1/5$</p>
